

Dear participant,

Thank you for your participation in nutritionDay worldwide in 2019 and for your effort. We are now able to present you your country report of the following sample size:

Number of participated center:	7
Number of participated units:	11
Number of patients:	113
Number of patients who gave consent:	113
Number of patients completing Sheet 3a:	113
Number of patients completing Sheet 3b:	112
Number of patients with 30-day outcome assessment:	96

This report compares your country data to international reference database based on data from nutritionDay 2016-2018.

The report consists of 4 parts:

Part I reflects capacity and staffing of your hospitals.

Part II reflects the organisation and the structure of the units.

Part III describes the patient's demographics, the clinical information and the outcome.

Part IV is the summary of information on patient's nutrition, history and health status.

Next year's nutritionDay will take place in November 2020. Again, we would very much appreciate your participation. You will be provided with further information on a regular

Best regards from the nutritionDay-Team!

Prof. Dr. Michael Hiesmayr

April Moicle

Sigrid Moick, MA



I. Hospital capacity and staffing ("Hospital sheet")

in hospital capacity and stanling (hospital sheet)		
	YOUR RESULTS	REFERENCE RESULTS
1. Total number of beds in hospital	152 [100-350]	333 [194-630]
2. Total number of admissions in the hospital last	8033 [4962-19214]	18005 [9101-36130]
year		
3. Total number of staff in the hospital		470 [00 000]
Total medical doctors	85 [49-107]	173 [80-399]
Medical specialists	56 [36-145]	116 [53-250]
Medical non-specialists	41 [38-74]	37 [14-84]
Nurses	316 [228-974]	420 [183-836]
Dieticians	4 [4-4]	5 [2-9]
Nutritionists	0 [0-1]	1 [0-4]
Pharmacists	34 [24-38]	7 [4-21]
Kitchen staff	28 [16-73]	31 [15-59]
Full time equivalent		
Total medical doctors	98 [98-98]	148 [60-373]
Medical specialists	51 [51-51]	98 [45-251]
Medical non-specialists	37 [37-37]	36 [11-99]
Nurses	222 [222-222]	364 [180-788]
Dieticians	1 [1-1]	5 [2-8]
Nutritionists	0 [0-0]	1 [0-3]
Pharmacists	16 [16-16]	6 [3-18]
Kitchen staff	8 [8-8]	29 [15-55]
4. Doos the bosnital have a nutrition care strategy?	4 (66 7%) Voc	408 (70 4%) Voc
4. Does the hospital have a nutrition care strategy?	4 (66,7%) Yes	498 (79,4%) Yes
5. Which nutrition-related standards or routine activities	s exist in your hospital?	
Nutrition training is available	5 (62,5%) Yes	434 (66,7%) Yes
Nutrition steering committee is available	1 (12,5%) Yes	388 (59,6%) Yes
Quality indicators are recorded and reported to national or regional level	3 (37,5%) Yes	263 (40,4%) Yes
Quality indicators are used for internal benchmarking	5 (62,5%) Yes	337 (51,8%) Yes
Patient feedback about food and food service is collected using a questionnaire	4 (50,0%) Yes	475 (73,0%) Yes
None	0 (0,0%) Yes	30 (4,6%) Yes
No answer given	2 (25,0%)	24 (3,7%)

6. Which codes are available /routinely used in your hospital for billing and reimbursement purposes?

	Codes	avai	lable
--	-------	------	-------

Nutrition Support

3 (37,5%) Yes

313 (48,1%) Yes





Oral nutrition supplements	4 (50,0%) Yes	239 (36,7%) Yes
Parenteral nutrition	2 (25,0%) Yes	375 (57,6%) Yes
Enteral nutrition	5 (62,5%) Yes	334 (51,3%) Yes
Dietary counseling	5 (62,5%) Yes	235 (36,1%) Yes
Specific dietary interventions	5 (62,5%) Yes	181 (27,8%) Yes
Screening for malnutrition	3 (37,5%) Yes	187 (28,7%) Yes
Risk of malnutrition	3 (37,5%) Yes	160 (24,6%) Yes
Malnutrition (in general)	4 (50,0%) Yes	294 (45,2%) Yes
Severity of malnutrition (i.e. mild, moderate, severe)	2 (25,0%) Yes	280 (43,0%) Yes
No information available from billing/finance/controlling	1 (12,5%) Yes	109 (16,7%) Yes
No answer given	-	42 (6,5%)
Codes routinely used		
Nutrition Support	3 (37,5%) Yes	270 (41,5%) Yes
Oral nutrition supplements	4 (50,0%) Yes	209 (32,1%) Yes
Parenteral nutrition	2 (25,0%) Yes	343 (52,7%) Yes
Enteral nutrition	5 (62,5%) Yes	307 (47,2%) Yes
Dietary counseling	5 (62,5%) Yes	198 (30,4%) Yes
Specific dietary interventions	5 (62,5%) Yes	159 (24,4%) Yes
Screening for malnutrition	4 (50,0%) Yes	149 (22,9%) Yes
Risk of malnutrition	2 (25,0%) Yes	128 (19,7%) Yes
Malnutrition (in general)	3 (37,5%) Yes	255 (39,2%) Yes
Severity of malnutrition (i.e. mild, moderate, severe)	1 (12,5%) Yes	242 (37,2%) Yes
No information available from billing/finance/controlling	1 (12,5%) Yes	120 (18,4%) Yes
No answer given	-	53 (8,1%)

II. Unit organisation and structures ("Sheet 1a/1b")

	YOUR RESULTS	REFERENCE RESULTS
Internal Medicine / General	4 (36,4%)	21.1%
Internal Medicine / Cardiology	0 (0,0%)	4.3%
Internal Medicine / Gastroenterology & hepatology	0 (0,0%)	7.8%
Internal Medicine / Geriatrics	0 (0,0%)	9.7%
Internal Medicine / Infectious diseases	0 (0,0%)	0.6%
Internal Medicine / Nephrology	0 (0,0%)	1.2%
Internal Medicine / Oncology (incl. radiotherapy)	0 (0,0%)	8.1%
Interdisciplinary	1 (9,1%)	3.4%
Long term care	0 (0,0%)	2.0%
Neurology	0 (0,0%)	3.6%
Surgery / General	2 (18,2%)	15.7%
Surgery/ Cardiac/Vascular/Thoracic	0 (0,0%)	1.4%
Surgery / Neurosurgery	0 (0,0%)	0.9%
Surgery / Orthopedic	0 (0,0%)	4.2%
Trauma	0 (0,0%)	0.9%
Ear Nose Throat (ENT)	0 (0,0%)	1.7%
Gynecology / Obstetrics	3 (27,3%)	1.4%
Pediatrics	0 (0,0%)	0.1%
Psychiatry	0 (0,0%)	0.9%
Others	1 (9,1%)	11.1%
2. Number of registered inpatients at noon	15 [10-25]	25 [19-32]
3. Total bed capacity of the unit	25 [19-28]	30 [24-40]
4. Number of each type of staff in the unit for TODAY's	morning shift	
Fully trained		
Medical doctors	3 [2-8]	4 [2-7]
Nurses	6 [4-9]	5 [3-8]
Nursing aides	1 [0-1]	2 [1-4]
Dieticians	1 [1-1]	1 [0-1]
Nutritionists	0 [0-1]	0 [0-1]
Administrative staff	1 [1-1]	1 [0-1]
Other staff involved in patient care	1 [1-1]	1 [0-2]
In training		
Medical doctors	0 [0-0]	1 [0-3]
Medical students	0 [0-4]	0 [0-2]

1 [0-3]

Nurses

0 [0-1]

Nursing aides	0 [0-0]	0 [0-0]
Dieticians	0 [0-0]	0 [0-0]
Nutritionists	0 [0-0]	0 [0-0]
Other staff involved in patient care	0 [0-0]	0 [0-0]
5. Is there a nutrition support team in your hospital available?	8 (100,0%) Yes	1018 (78,3%) Yes
6. Does the unit have a nutrition care strategy?	8 (100,0%) Yes	984 (75,7%) Yes
7. Is there a person in your unit responsible for nutrition care?	7 (87,5%) Yes	922 (70,9%) Yes
8. Is there a dietician, nutritionist or dietetic assistant available for your unit?	10 (100,0%) Yes	1239 (90,8%) Yes
9. Is specific staff responsible for providing feeding assistance to patients during meal times?	3 (37,5%) Yes	780 (60,0%) Yes
10. How do you MAINLY screen/monitor patients for malnutrition?		
At admission		
No routine screening	3 (27,3%) Yes	109 (7,6%)
No fixed criteria	-	21 (1,5%)
Experience / visual assessment only	-	102 (7,1%)
Weighing / BMI only	7 (63,6%) Yes	201 (14,1%)
Nutritional Risk Screening (NRS) 2002	-	462 (32,3%)
Malnutrition Universal Screening Tool (MUST)	-	81 (5,7%)
Malnutrition Screening tool (MST)	-	115 (8,0%)
SNAQ	-	13 (0,91%)
Other formal tool	-	249 (17,4%)
I do not know	-	11 (0,77%)
Missing	1 (9,1%)	66 (4,6%)
During hospital stay		
No routine monitoring	2 (18,2%) Yes	133 (9,3%)
No fixed criteria	1 (9,1%) Yes	118 (8,3%)
Experience / visual assessment only	2 (18,2%) Yes	235 (16,4%)
Weighing / BMI only	4 (36,4%) Yes	471 (32,9%)
Other formal tool	1 (9,1%) Yes	375 (26,2%)
I do not know	-	32 (2,2%)
Missing	1 (9,1%)	66 (4,6%)
11a. Do you routinely use guidelines or standards for nutrition care?	8 (100,0%) Yes	1043 (82,1%) Yes

11b. If yes, which one is mainly used?		
International guidelines	2 (25,0%) Yes	257 (24,6%)
National guidelines	-	166 (15,9%)
Standards on hospital level	3 (37,5%) Yes	378 (36,2%)
Standards on unit level	1 (12,5%) Yes	65 (6,2%)
Individual patient nutrition care plans	2 (25,0%) Yes	155 (14,9%)
Other	-	11 (1,1%)
Missing	-	11 (1,1%)

12. What is routinely done in your unit for given patient groups?

At risk		
Watchful waiting	3 (27,3%) Yes	399 (27,9%)
Discuss nutrition care activities during ward rounds	5 (45,5%) Yes	614 (42,9%)
Develop an individual nutrition care plan	4 (36,4%) Yes	720 (50,3%)
Initiate treatment / nutrition intervention	4 (36,4%) Yes	820 (57,3%)
Consult a nutrition expert (dietician, nutritionist, etc.)	4 (36,4%) Yes	828 (57,9%)
Consult a medical professional	4 (36,4%) Yes	505 (35,3%)
Calculate energy requirements	3 (27,3%) Yes	652 (45,6%)
Calculate protein requirements	6 (54,5%) Yes	630 (44,1%)
Malnourished		
Watchful waiting	1 (9,1%) Yes	280 (19,6%)
Discuss nutrition care activities during ward rounds	1 (9,1%) Yes	675 (47,2%)
Develop an individual nutrition care plan	1 (9,1%) Yes	792 (55,4%)
Initiate treatment / nutrition intervention	2 (18,2%) Yes	884 (61,8%)
Consult a nutrition expert (dietician, nutritionist, etc.)	2 (18,2%) Yes	823 (57,6%)
Consult a medical professional	-	579 (40,5%)
Calculate energy requirements	5 (45,5%) Yes	763 (53,4%)
Calculate protein requirements	3 (27,3%) Yes	745 (52,1%)
Every patient		
Watchful waiting	3 (27,3%) Yes	700 (49,0%)
Discuss nutrition care activities during ward rounds	3 (27,3%) Yes	358 (25,0%)
Develop an individual nutrition care plan	4 (36,4%) Yes	250 (17,5%)
Initiate treatment / nutrition intervention	3 (27,3%) Yes	197 (13,8%)
Consult a nutrition expert (dietician, nutritionist, etc.)	1 (9,1%) Yes	264 (18,5%)
Consult a medical professional	2 (18,2%) Yes	307 (21,5%)
Calculate energy requirements	1 (9,1%) Yes	186 (13,0%)
Calculate protein requirements	-	168 (11,7%)
Never		

Watchful waiting

2 (18,2%) Yes

105 (7,3%)



Discuss nutrition care activities during ward rounds	<u>.</u>	77 (5,4%)
Develop an individual nutrition care plan	<u> </u>	62 (4,3%)
Initiate treatment / nutrition intervention	-	19 (1,3%)
Consult a nutrition expert (dietician, nutritionist, etc.)	2 (18,2%) Yes	20 (1,4%)
Consult a medical professional	2 (18,2%) Yes	146 (10,2%)
Calculate energy requirements	-	141 (9,9%)
Calculate protein requirements		165 (11,5%)
		100 (11,070)
l do not know		
Watchful waiting	-	71 (5,0%)
Discuss nutrition care activities during ward rounds	1 (9,1%) Yes	54 (3,8%)
Develop an individual nutrition care plan	-	41 (2,9%)
Initiate treatment / nutrition intervention	-	34 (2,4%)
Consult a nutrition expert (dietician, nutritionist, etc.)	-	33 (2,3%)
Consult a medical professional	-	86 (6,0%)
Calculate energy requirements		65 (4,5%)
Calculate protein requirements	1 (9,1%) Yes	71 (5,0%)
		(, ,
13. When do you routinely weigh your patients?		
at admission	8 (72,7%) Yes	928 (64,9%)
Within 24 hours	-	210 (14,7%)
Within 48 hours	-	79 (5,5%)
Within 72 hours	1 (9,1%) Yes	39 (2,7%)
Every week	-	627 (43,8%)
Occasionally	-	158 (11,0%)
When requested	7 (63,6%) Yes	728 (50,9%)
At discharge	-	67 (4,7%)
Never	-	17 (1,2%)
l do not know	-	6 (0,42%)
No answer given	3 (16,7%)	69 (4,8%)
14. What do you do to support adequate food intake of p	patients?	
Offer additional meals or in between snacks	6 (54,5%) Yes	1096 (76,6%)
Offer meal choices	8 (72,7%) Yes	1055 (73,8%)
Offer different portion sizes	1 (9,1%) Yes	934 (65,3%)
Consider food presentation	5 (45,5%) Yes	502 (35,1%)
Change food texture/consistency as needed	5 (45,5%) Yes	1175 (82,2%)
Consider patient problems with eating and drinking	5 (45,5%) Yes	1166 (81,5%)
Ensure that mealtimes are undisturbed/protected mealtime policy	6 (54,5%) Yes	324 (22,7%)
Promote positive eating environment	5 (45,5%) Yes	419 (29,3%)
Consider cultural/religious preferences	5 (45,5%) Yes	903 (63,1%)
Consider patient allergies / intolerances	7 (63,6%) Yes	1045 (73,1%)
Other	-	103 (7,2%)



		26 (4.99()
I do not know	-	26 (1,8%)
No answer given	1 (9,1%)	67 (4,7%
15. Which nutrition-related standards or routine activities	exist in your unit?	
Nutrition training is available	2 (18,2%) Yes	821 (57,4%)
Reporting of nutrition related information to hospital managers	4 (36,4%) Yes	608 (42,5%)
Quality indicators are recorded and reported to national or regional level	3 (27,3%) Yes	448 (31,3%)
Quality indicators are used for internal benchmarking	4 (36,4%) Yes	513 (35,9%
Patient feedback about food and food service is collected using a questionnaire	6 (54,5%) Yes	884 (61,8%)
None	-	41 (2,9%)
I do not know	-	61 (4,3%)
No answer given	3 (27,3%)	151 (10,6%)
40. At admission what is called and decomposited		
16. At admission what is asked and documented?	E(AEEQ/) Vac	1104 (77 20/
Change in weight	5 (45,5%) Yes	1104 (77,2%)
Eating habits/difficulties	5 (45,5%) Yes	1041 (72,8%)
Nutrition before admission	1 (9,1%) Yes	823 (57,6%
None	-	18 (1,3%
I do not know	-	33 (2,3%
No answer given	3 (27,3%)	153 (10,7%)
17. On what forms is there a specific part about eating, nu	trition or malnutrition?	
a. Patient Record has a section for		
indicating if the patient is malnourished or at risk of malnutrition	3 (27,3%) Yes	1037 (72,5%)
nutrition treatment	4 (36,4%) Yes	837 (58,5%)
None	2 (18,2%) Yes	65 (4,5%)
I do not know	-	42 (2,9%)
b. Discharge Letter		
summarizes nutrition treatment received during stay	2 (18,2%) Yes	592 (41,4%)
makes future nutrition-related recommendations	5 (45,5%) Yes	816 (57,1%)
None	2 (18,2%) Yes	137 (9,6%)
I do not know	2 (10,270) 165	128 (9,0%
	-	128 (9,0 %
18. Do you provide brochures about malnutrition to at risk/malnourished patients?	3 (37,5%) Yes	519 (45,2%) Yes
19. Who filled in this sheet?		
Head staff	1 (9,1%) Yes	411 (28,7%)
Dietician	4 (36,4%) Yes	717 (50,1%)
		• •

388 (27,1%)

Nurse

4 (36,4%) Yes

Physician	-	175 (12,2%)
Administrative staff	-	13 (0,91%)
Other	-	71 (5,0%)
None	-	-
l do not know	-	15 (1,0%)



	YOUR RESULTS	REFERENCE RESULTS
Total	113	27195
Age	40 [30-64]	67 [53-78]
Female	83 (73,5%)	13525 (49,7%)
Weight	69,8±19,6	69,9±18,9
Height	159±7	165±10
BMI	27,4±7,6	25,5±5,9
1. This hospital admission was…		
planned	50 (44,2%)	9881 (36,3%)
an emergency	57 (50,4%)	14977 (55,1%)
I do not know	6 (5,3%)	2337 (8,6%)
No answer given	-	-
2a. Diagnosis at admission		
0100 Infectious and parasitic diseases	2 (1,8%)	2442 (9,0%)
0200 Neoplasms	7 (6,2%)	4873 (17,9%)
0300 Blood and bloodforming organs and the immune mechanism	7 (6,2%)	1724 (6,3%)
0400 Endocrine, nutritional and metabolic diseases	8 (7,1%)	3266 (12,0%)
0500 Mental health	2 (1,8%)	1218 (4,5%)
0600 Nervous system	2 (1,8%)	2493 (9,2%)
0700 Eye and adnexa	1 (0,88%)	339 (1,2%)
0800 Ear and mastoid process	1 (0,88%)	172 (0,63%)
0900 Circulatory system	9 (8,0%)	5629 (20,7%)
1000 Respiratory system	18 (15,9%)	4217 (15,5%)
1100 Digestive system	21 (18,6%)	6840 (25,2%)
1200 Skin and subcutaneous tissue	8 (7,1%)	1244 (4,6%)
1300 Musculoskeletal system and connective tissue	4 (3,5%)	4246 (15,6%)
1400 Genitourinary system	6 (5,3%)	2809 (10,3%)
1500 Pregnancy, childbirth and the puerperium	30 (26,5%)	222 (0,82%)
1600 Conditions originating in the perinatal period	2 (1,8%)	36 (0,13%)
1700 Congenital/chromosomal abnormalities	-	52 (0,19%)
1800 Symptoms, signs, abnormal clinical/lab findings	5 (4,4%)	1319 (4,9%)
1900 Injury, poisoning	-	724 (2,7%)
2000 External causes of morbidity and mortality (e.g. transport accidents, assaults)	1 (0,88%)	671 (2,5%)
2100 Factors influencing health status and contact with health services	11 (9,7%)	990 (3,6%)
No answer given	-	-

3. Which conditions/comorbidities does this patient have?



Myocardial infarction	5 (4,5%)	1338 (5,5%)
Chronic lung disease	6 (5,4%)	3466 (14,1%)
Cerebral vascular disease	-	2274 (9,3%)
Peripheral vascular disease	2 (1,8%)	2842 (11,6%)
Chronic liver disease	2 (1,8%)	1477 (6,1%)
Chronic kidney disease	5 (4,5%)	2817 (11,6%)
Diabetes	28 (25,2%)	5968 (24,2%)
Cancer	17 (15,3%)	5709 (23,2%)
Infection	9 (8,1%)	3578 (14,6%)
Dementia	1 (0,90%)	1229 (5,1%)
Major depressive disorder	-	1233 (5,1%)
Other chronic mental disorder	2 (1,8%)	1229 (5,1%)
Other chronic disease	14 (12,6%)	6369 (25,9%)
None	51 (45,1%)	4869 (17,9%)
4a. Previous operation during this hospital stay		
Yes, planned	10 (8,8%)	5304 (19,5%)
Yes, acute	-	1634 (6,0%)
No	79 (69,9%)	17917 (65,9%)
I do not know	2 (1,8%)	313 (1,2%)
Missing	22 (19,5%)	2027 (7,5%)
Days since operation	2 [2-2]	3 [1-11]
4b. Planned operation during this hospital stay	10 (0.00()	4000 (7.00()
Yes, today or tomorrow	10 (8,8%)	1998 (7,3%)
Yes, later	2 (1,8%)	1411 (5,2%)
No	76 (67,3%)	19908 (73,2%)
I do not know	2 (1,8%)	1082 (4,0%)
Missing	23 (20,4%)	2796 (10,3%)
5. Previous ICU admission during this hospital stay?	2 (2,2%)	2731 (10,6%)
$\mathbf{J}_{\mathbf{i}}$	2 (2,270)	2781 (10,070)
(Yes)		
	24 (21,2%)	1729 (6,4%)
(Yes)	24 (21,2%)	1729 (6,4%)
(Yes)	24 (21,2%)	1729 (6,4%)
(Yes) 6. Is this patient terminally ill?	24 (21,2%) 65 (57,5%)	1729 (6,4%) 20820 (76,6%)
(Yes) 6. Is this patient terminally ill? 7. Fluid status		
(Yes) 6. Is this patient terminally ill? 7. Fluid status Normal	65 (57,5%)	20820 (76,6%) 1466 (5,4%) 1298 (4,8%)
(Yes) 6. Is this patient terminally ill? 7. Fluid status Normal Overloaded	65 (57,5%) 29 (25,7%)	20820 (76,6%) 1466 (5,4%)
(Yes) 6. Is this patient terminally ill? 7. Fluid status Normal Overloaded Dehydrated	65 (57,5%) 29 (25,7%) 14 (12,4%)	20820 (76,6%) 1466 (5,4%) 1298 (4,8%)



At risk 9 (8,0%) 4856 (17,9%) No 76 (67,3%) 16820 (61,8%) I do not know 10 (8,8%) 2255 (8,3%) Missing - - 10. IV Fluids - - Electrolyte solution (NaCl, Ringers lactate, etc) 10 (8,8%) 9496 (34,9%) 5% Glucose solution 16 (14,2%) 2615 (9,6%) 11. Number of ONS drinks planned 0 [0-2] 0 [0-0] 12. Nutrition intake - - Regular hospital food 93 (82,3%) 15788 (58,1%) Fortlifie/drin/tiched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) 13a. All lines and Tubes - 2360 (9,2%) Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogastric 2 (2,2%) 680 (2,6%)	9 Number of different mediactions related		
Other 2 [1-2] 2 [1-4] 9. Was this patient identified as malnourished or at risk of malnutrition? Malnourished 18 (15,9%) 3264 (12,0%) At risk 9 (8,0%) 4856 (17,9%) No 76 (67,3%) 16820 (61,8%) 1 do not know 10 (8,8%) 2255 (8,3%) 16820 (61,8%) 10 (8,8%) 2255 (8,3%) 10. IV Fluids Electrolyte solution (NaCl, Ringers lactate, etc) 10 (8,8%) 9496 (34,9%) 5% Glucose solution 16 (14,2%) 2615 (9,6%) 11. Number of ONS drinks planned 0 [0-2] 0 [0-0] 0 [0-0] 12. Nutrition intake Regular hospital food 93 (82,3%) 15788 (58,1%) Fortified/enriched hospital food 93 (82,3%) 15788 (58,1%) 101 (14,8%) Parenteral nutrition 6 (5,3%) 1315 (4,8%) 126 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) 1283 (4,7%) None 3 (2,7%) 1283 (4,7%) 1283 (4,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogastric 2		0 [4 0]	E [0, 0]
9. Was this patient identified as malnourished or at risk of malnutrition? Malnourished 18 (15,9%) 3264 (12,0%) At risk 9 (8,0%) 4456 (17,9%) No 76 (67,3%) 16820 (61,8%) I do not know 10 (8,8%) 2255 (8,3%) Missing - - 10. IV Fluids - - Electrolyte solution (NaCl, Ringers lactate, etc) 10 (8,8%) 9496 (34,9%) 5% Glucose solution 16 (14,2%) 2615 (9,6%) 11. Number of ONS drinks planned 0 [0-2] 0 [0-0] 12. Nutrition intake - - Regular hospital food 93 (82,3%) 15788 (58,1%) Fortified/enriched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Parenteral nutrition 4 (3,5%) 1286 (4,7%) None 3 (2,7%) 1283 (4,7%) 13. All lines and Tubes - - Central Venous 2 (2,2%) 680 (2,6%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 136 (0,63%)			
Malnourished 18 (15,9%) 3264 (12,0%) At risk 9 (8,0%) 4456 (77,9%) No 76 (67,3%) 16820 (61,8%) I on to know 10 (8,8%) 2255 (8,3%) Missing - - 10. IV Fluids - - Electrolyte solution (NaCl, Ringers lactate, etc) 10 (8,8%) 9496 (34,9%) 5% Glucose solution 16 (14,2%) 2615 (9,6%) 11. Number of ONS drinks planned 0 [0-2] 0 [0-0] 12. Nutrition intake - - Regular hospital food 93 (82,3%) 15788 (58,1%) Fortified/enriched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Enteral nutrition 4 (3,5%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogastric 2 34 (0,91%) 126 (0,499%)	Other	2 [1-2]	2 [1-4]
At risk 9 (8,0%) 4456 (17,9%) No 76 (67,3%) 16820 (61,8%) 10 (8,8%) 2255 (8,3%) Missing - - - 10. IV Fluids - - - Electrolyte solution (NaCl, Ringers lactate, etc) 10 (8,8%) 9496 (34,9%) 5% Glucose solution 16 (14,2%) 2615 (9,6%) 11. Number of ONS drinks planned 0 [0-2] 0 [0-0] 12. Nutrition intake - - Regular hospital food 93 (82,3%) 15788 (58,1%) Fortfied/enriched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) 13a. All lines and Tubes - 2360 (9,2%) Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasoduadenal 3 (3,3%) 99 (0,39%) Peroutaneous endoscopy/surgical gastrostomy -	9. Was this patient identified as malnourished or at risk of	f malnutrition?	
No 76 (67,3%) 16820 (61,8%) I do not know 10 (8,8%) 2255 (8,3%) Missing - - 10. IV Fluids - - Electrolyte solution (NaCl, Ringers lactate, etc) 10 (8,8%) 9496 (34,9%) 5% Glucose solution 16 (14,2%) 2615 (9,6%) 11. Number of ONS drinks planned 0 [0-2] 0 [0-0] 12. Nutrition intake - - Regular hospital food 93 (82,3%) 117588 (58,1%) Fortified/enriched hospital food 4 (3,5%) 4012 (14,8%) Enteral nutrition 6 (5,3%) 4012 (14,8%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 115 (4,8%) Parenteral nutrition 4 (3,5%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None - 2360 (9,2%) Peripheral venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasoduadenal 3 (3,3%) 99 (0,39%) Peroutaneous endoscopy/surgical gastrostomy <t< td=""><td>Malnourished</td><td>18 (15,9%)</td><td>3264 (12,0%)</td></t<>	Malnourished	18 (15,9%)	3264 (12,0%)
l do not know 10 (8,8%) 2255 (8,3%) Missing	At risk	9 (8,0%)	4856 (17,9%)
Missing - - 10. IV Fluids - - Electrolyte solution (NaCl, Ringers lactate, etc) 10 (8,8%) 9496 (34,9%) 5% Glucose solution 16 (14,2%) 2615 (9,6%) 11. Number of ONS drinks planned 0 [0-2] 0 [0-0] 12. Nutrition intake - - Regular hospital food 93 (82,3%) 15788 (58,1%) Fortified/enriched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Parenteral nutrition 6 (5,3%) 1286 (4,7%) Special diet 14 (12,4%) 8824 (32,5%) None 3 (2,7%) 1283 (4,7%) 13. All lines and Tubes - 2360 (9,2%) Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogastric 2 (3,0%) 99 (0,39%) Percutaneous endoscopy/surgical gastrostomy - 126 (0,49%) Percutaneous endoscopy/surgical gas	No	76 (67,3%)	16820 (61,8%)
10. IV Fluids Electrolyte solution (NaCl, Ringers lactate, etc) 10 (8,8%) 9496 (34,9%) 5% Glucose solution 16 (14,2%) 2615 (9,6%) 11. Number of ONS drinks planned 0 [0-2] 0 [0-0] 12. Nutrition intake 15788 (58,1%) 5768 (58,1%) Fortified/enriched hospital food 93 (82,3%) 15788 (58,1%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 1315 (4,8%) Parenteral nutrition 4 (3,5%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) 13. All lines and Tubes 2 2,2%) Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasoglastric 2 (2,2%) 680 (2,6%) Nasoglastric 2 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy 95 (0,37%) 1200 (48,5%) <td>I do not know</td> <td>10 (8,8%)</td> <td>2255 (8,3%)</td>	I do not know	10 (8,8%)	2255 (8,3%)
Electrolyte solution (NaCl, Ringers lactate, etc) 10 (8,8%) 9496 (34,9%) 5% Glucose solution 16 (14,2%) 2615 (9,6%) 11. Number of ONS drinks planned 0 [0-2] 0 [0-0] 12. Nutrition intake 7 7 Regular hospital food 93 (82,3%) 15788 (58,1%) Fortified/enriched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Parenteral nutrition 6 (5,3%) 1315 (4,8%) Parenteral nutrition 6 (5,3%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) None 3 (2,7%) 1280 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogiginal 2 (2,2%) 680 (2,6%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 362 (1,3%) <td>Missing</td> <td>-</td> <td>-</td>	Missing	-	-
Electrolyte solution (NaCl, Ringers lactate, etc) 10 (8,8%) 9496 (34,9%) 5% Glucose solution 16 (14,2%) 2615 (9,6%) 11. Number of ONS drinks planned 0 [0-2] 0 [0-0] 12. Nutrition intake 7 7 Regular hospital food 93 (82,3%) 15788 (58,1%) Fortified/enriched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Parenteral nutrition 6 (5,3%) 1315 (4,8%) Parenteral nutrition 6 (5,3%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) Parenteral nutrition 2 (2,2%) 6680 (2,6%) Nasogiginal 2 (2,2%) 6680 (2,6%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma 126 (0,49%) 1280 (45,5%) Percutaneous endoscopy/surgical gastrostomy 2 324 (0,91%) 12303 (45,2%) None 1200 (88,5%) 12303 (45,2%) 12303 (45,2%) None 1200 (88,5%)	10. IV Fluids		
5% Glucose solution 16 (14,2%) 2615 (9,6%) 11. Number of ONS drinks planned 0 [0-2] 0 [0-0] 12. Nutrition intake 8 15788 (58,1%) Fortified/enriched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Parenteral nutrition 6 (5,3%) 4012 (14,8%) Parenteral nutrition 6 (5,3%) 1315 (4,8%) Parenteral nutrition 4 (3,5%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) 13a. All lines and Tubes 2 2,2%) Central Venous - 2360 (9,2%) Preipheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogastric 2 (2,2%) 680 (2,6%) Nasoduadenal 3 (3,3%) 99 (0,39%) Percutaneous endoscopy/surgical gastrostomy - 323 (0,91%) Percutaneous endoscopy/surgical gastrostomy - 35 (0,37%) None 100 (88,5%) 12303 (45,2%) 13b.		10 (8.8%)	9496 (34.9%)
11. Number of ONS drinks planned 0 [0-2] 0 [0-0] 12. Nutrition intake 93 (82,3%) 15788 (58,1%) Regular hospital food 93 (82,3%) 15788 (58,1%) Fortified/enriched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Parenteral nutrition 6 (5,3%) 1315 (4,8%) Parenteral nutrition 4 (3,5%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) Task All lines and Tubes Central Venous Central Venous 2 (2,2%) Central Venous 2 (2,2%) 680 (2,6%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogastric 2 (2,2%) 680 (2,6%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 135 (0,53%) Nasoduadenal 3 (3,3%) 99 (0,37%) None 100 (88,5%) 12303 (45,2%) Task All lines and Tubes - Task Contraneous endoscopy/surgical gastro			· · · · · ·
12. Nutrition intake Regular hospital food 93 (82,3%) 15788 (58,1%) Fortified/enriched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Enteral nutrition 6 (5,3%) 4012 (14,8%) Parenteral nutrition 6 (5,3%) 1315 (4,8%) Parenteral nutrition 4 (3,5%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) 13a. All lines and Tubes - 2360 (9,2%) Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogiaptinal - 135 (0,53%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) 13b. Were there complications with nutrition related lines and tubes s			(,,,,,)
Regular hospital food 93 (82,3%) 15788 (58,1%) Fortified/enriched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Enteral nutrition 6 (5,3%) 1315 (4,8%) Parenteral nutrition 4 (3,5%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) Tas. All lines and Tubes - Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogastric 2 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) Tab. Were there complications with nutrition related lines and tubes since admission? 3 (2,7%) 227 (0,83%) Yes, ongoing 3 (2,7%) 21227 (78,1%) <td>11. Number of ONS drinks planned</td> <td>0 [0-2]</td> <td>0 [0-0]</td>	11. Number of ONS drinks planned	0 [0-2]	0 [0-0]
Regular hospital food 93 (82,3%) 15788 (58,1%) Fortified/enriched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Enteral nutrition 6 (5,3%) 1315 (4,8%) Parenteral nutrition 4 (3,5%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) Tas. All lines and Tubes - Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogastric 2 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) Tab. Were there complications with nutrition related lines and tubes since admission? 3 (2,7%) 227 (0,83%) Yes, ongoing 3 (2,7%) 227 (0,83%)	12. Nutrition intake		
Fortified/enriched hospital food 4 (3,5%) 3133 (11,5%) Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Enteral nutrition 6 (5,3%) 1315 (4,8%) Parenteral nutrition 4 (3,5%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) Total Venous 3 (2,7%) Total Venous - Central Venous - Central Venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogiejunal - 135 (0,53%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical gastrostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) State of the proving the nutrition related lines and tubes since admission? Yes, ongoing 3 (2		93 (82.3%)	15788 (58.1%)
Protein/energy supplement (e.g. ONS drinks) 6 (5,3%) 4012 (14,8%) Enteral nutrition 6 (5,3%) 1315 (4,8%) Parenteral nutrition 4 (3,5%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) 13a. All lines and Tubes - 2360 (9,2%) Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogiejunal - 135 (0,53%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) 13b. Were there complications with nutrition related lines and tubes since admission? - 362 (1,3%) Yes, previously - 362 (1,3%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) 10	•		, , , , , , , , , , , , , , , , , , ,
Enteral nutrition 6 (5,3%) 1315 (4,8%) Parenteral nutrition 4 (3,5%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) 13a. All lines and Tubes 3 (2,7%) 1283 (4,7%) Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasogiaptinal - 135 (0,53%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) 13b. Were there complications with nutrition related lines and tubes since admission? - 362 (1,3%) Yes, previously - 362 (1,3%) Yes, ongoing 3 (2,7%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) I do not know 7 (6,2	-	, · · · ,	, , , , , , , , , , , , , , , , , , ,
Parenteral nutrition 4 (3,5%) 1286 (4,7%) Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) Table State Stat			
Special diet 14 (12,4%) 8834 (32,5%) None 3 (2,7%) 1283 (4,7%) 13a. All lines and Tubes Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasojejunal - 135 (0,53%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) 13b. Were there complications with nutrition related lines and tubes since admission? - 362 (1,3%) Yes, previously - 362 (1,3%) 3 (2,7%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) 1d on tknow 7 (6,2%) 2312 (8,5%)			,
None 3 (2,7%) 1283 (4,7%) 13a. All lines and Tubes - 2360 (9,2%) Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasojejunal - 135 (0,53%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) T - 362 (1,3%) Yes, previously - 362 (1,3%) Yes, ongoing 3 (2,7%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) I do not know 7 (6,2%) 2312 (8,5%)			
Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasojejunal - 135 (0,53%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) 13b. Were there complications with nutrition related lines and tubes since admission? - 362 (1,3%) Yes, previously - 362 (1,3%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) 1 I do not know 7 (6,2%) 2312 (8,5%) 1	None		
Central Venous - 2360 (9,2%) Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasojejunal - 135 (0,53%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) 13b. Were there complications with nutrition related lines and tubes since admission? - 362 (1,3%) Yes, previously - 362 (1,3%) Yes, ongoing 3 (2,7%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) I do not know 7 (6,2%) 2312 (8,5%)			
Peripheral venous access 10 (11,0%) 12501 (48,7%) Nasogastric 2 (2,2%) 680 (2,6%) Nasojejunal - 135 (0,53%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) 13b. Were there complications with nutrition related lines and tubes since admission? - 362 (1,3%) Yes, ongoing 3 (2,7%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) I do not know 7 (6,2%) 2312 (8,5%)			
Nasogastric 2 (2,2%) 680 (2,6%) Nasojejunal - 135 (0,53%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) - - 362 (1,3%) Yes, ongoing 3 (2,7%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) I do not know 7 (6,2%) 2312 (8,5%)		-	
Nasojejunal - 135 (0,53%) Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) - - 362 (1,3%) Yes, previously - 362 (1,3%) Yes, ongoing 3 (2,7%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) I do not know 7 (6,2%) 2312 (8,5%)	-		
Nasoduadenal 3 (3,3%) 99 (0,39%) Enterostoma - 126 (0,49%) Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) 13b. Were there complications with nutrition related lines and tubes since admission? - 362 (1,3%) Yes, previously - 362 (1,3%) No 57 (50,4%) 21227 (78,1%) I do not know 7 (6,2%) 2312 (8,5%)		2 (2,2%)	,
Enterostoma - 126 (0,49%) 126 (0,49%) 126 (0,49%) 126 (0,49%) 126 (0,49%) 126 (0,49%) 1234 (0,91%) 234 (0,91%) 234 (0,91%) 234 (0,91%) 12303 (45,2%) 12303 (45,		-	
Percutaneous endoscopy/surgical gastrostomy - 234 (0,91%) Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) 13b. Were there complications with nutrition related lines and tubes since admission? - 362 (1,3%) Yes, previously - 362 (1,3%) Yes, ongoing 3 (2,7%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) I do not know 7 (6,2%) 2312 (8,5%)		3 (3,3%)	,
Percutaneous endoscopy/surgical jejunostomy - 95 (0,37%) None 100 (88,5%) 12303 (45,2%) 13b. Were there complications with nutrition related lines and tubes since admission? - 362 (1,3%) Yes, previously - 362 (1,3%) Yes, ongoing 3 (2,7%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) I do not know 7 (6,2%) 2312 (8,5%)		-	
None 100 (88,5%) 12303 (45,2%) 13b. Were there complications with nutrition related lines and tubes since admission? - - Yes, previously - 362 (1,3%) Yes, ongoing 3 (2,7%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) I do not know 7 (6,2%) 2312 (8,5%)	Percutaneous endoscopy/surgical gastrostomy	-	· · · · · · · · · · · · · · · · · · ·
13b. Were there complications with nutrition related lines and tubes since admission? Yes, previously-362 (1,3%)Yes, ongoing3 (2,7%)227 (0,83%)No57 (50,4%)21227 (78,1%)I do not know7 (6,2%)2312 (8,5%)	Percutaneous endoscopy/surgical jejunostomy	-	95 (0,37%)
and tubes since admission? Yes, previously - 362 (1,3%) Yes, ongoing 3 (2,7%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) I do not know 7 (6,2%) 2312 (8,5%)	None	100 (88,5%)	12303 (45,2%)
Yes, ongoing 3 (2,7%) 227 (0,83%) No 57 (50,4%) 21227 (78,1%) I do not know 7 (6,2%) 2312 (8,5%)	13b. Were there complications with nutrition related lines and tubes since admission?		
No57 (50,4%)21227 (78,1%)I do not know7 (6,2%)2312 (8,5%)	Yes, previously	-	362 (1,3%)
I do not know 7 (6,2%) 2312 (8,5%)	Yes, ongoing	3 (2,7%)	227 (0,83%)
	No	57 (50,4%)	21227 (78,1%)
Missing 46 (40,7%) 3067 (11,3%)	I do not know	7 (6,2%)	2312 (8,5%)
	Missing	46 (40,7%)	3067 (11,3%)



14. Please indicate if any of the following was done for this	s nationt since admissio	n
Energy requirements were determined	22 (24,2%)	9593 (37,3%)
Protein requirements were determined	9 (9,9%)	8543 (33,3%)
•		
Food/Nutrition intake was recorded in the patient record	16 (17,6%)	11079 (43,1%)
Nutrition treatment plan was developed	12 (13,2%)	9042 (35,2%)
Nutrition expert was consulted	11 (12,1%)	9313 (36,3%)
Malnutrition status is recorded in the patient record	9 (9,9%)	8238 (32,1%)
None	-	-
15a. Energy goal		
< 500 kcal	19 (16,8%)	1046 (3,8%)
500-999 kcal	3 (2,7%)	269 (0,99%)
1000-1499 kcal	10 (8,8%)	2400 (8,8%)
1500-1999 kcal	34 (30,1%)	8218 (30,2%)
>=2000 kcal	2 (1,8%)	2704 (9,9%)
Not determined	16 (14,2%)	8749 (32,2%)
I do not know	7 (6,2%)	2265 (8,3%)
Missing	22 (19,5%)	1544 (5,7%)
J. J		
15b. Energy intake		
< 500 kcal	19 (16,8%)	1697 (6,2%)
500-999 kcal	8 (7,1%)	1257 (4,6%)
1000-1499 kcal	12 (10,6%)	3424 (12,6%)
1500-1999 kcal	29 (25,7%)	5999 (22,1%)
>=2000 kcal	2 (1,8%)	1448 (5,3%)
Not determined	15 (13,3%)	8533 (31,4%)
I do not know	6 (5,3%)	3276 (12,0%)
Missing	22 (19,5%)	1561 (5,7%)
16. Since admission, this patient's health status has		
Improved	44 (38,9%)	12837 (47,2%)
Deteriorated	6 (5,3%)	1433 (5,3%)
Remained the same	28 (24,8%)	7057 (25,9%)
This patient has just been admitted	4 (3,5%)	1699 (6,2%)
I do not know	9 (8,0%)	2663 (9,8%)
Missing	22 (19,5%)	1506 (5,5%)
Length of hospital stay (days)	6 [4-13]	12 [6-23]
	0[110]	.2 [0 20]
Outcome Code		
1= Still in the hospital	20 (17,7%)	2601 (9,6%)
2= Transferred to another hospital	1 (0,88%)	571 (2,1%)
3= Transferred to long term care	-	1107 (4,1%)

1 (0,88%)	1222 (4,5%)
71 (62,8%)	20149 (74,1%)
2 (1,8%)	845 (3,1%)
1 (0,88%)	507 (1,9%)
17 (15,0%)	193 (0,71%)
53 (70,7%)	18115 (75,8%)
9 (12,0%)	1637 (6,9%)
2 (2,7%)	1465 (6,1%)
-	112 (0,47%)
-	112 (0,47%)
1 (1,3%)	796 (3,3%)
10 (13,3%)	1597 (6,7%)
	71 (62,8%) 2 (1,8%) 1 (0,88%) 17 (15,0%) 53 (70,7%) 9 (12,0%) 2 (2,7%) - - 1 (1,3%)



	YOUR RESULTS	REFERENCE RESULTS
1. What are your typical dietary habits?		
No special dietary habits	75 (66,4%)	17526 (66,5%)
I am vegetarian	4 (3,5%)	510 (1,9%)
I adhee to a vegan diet	4 (3,5%)	193 (0,73%)
I eat gluten-free diet	3 (2,7%)	244 (0,93%
I avoid added sugars	17 (15,0%)	3934 (14,9%)
I avoid carbohydrates	1 (0,88%)	1502 (5,7%
I eat a low fat-diet	5 (4,4%)	2899 (11,0%)
I am lactose intolerant	-	946 (3,6%)
Other special diet due to intolerances/allergies	2 (1,8%)	469 (1,8%)
Other	5 (4,4%)	1854 (7,0%)
No answer given	24 (21,2%)	1891 (7,2%)
2. Where did you live before your current hospital admini-	ssion?	
At home	90 (79,6%)	22025 (83,6%)
In a nursing home or other live-in facility	-	836 (3,2%)
I was transferred from another hospital	1 (0,88%)	1411 (5,4%)
Other	-	374 (1,4%)
Missing	22 (19,5%)	1694 (6,4%)
3. In general, are you able to walk?		
Yes	68 (60,2%)	16014 (60,8%)
Yes, with someone's help	3 (2,7%)	2580 (9,8%)
Yes, independently using a cane, walker, or crutches	7 (6,2%)	3466 (13,2%)
No, I have a wheelchair	8 (7,1%)	1033 (3,9%)
No, I am bedridden	4 (3,5%)	1462 (5,6%)
Missing	23 (20,4%)	1785 (6,8%)
5		
4. In general, how would you say your health is?		
Very good	30 (26,5%)	1854 (7,0%)
Good	28 (24,8%)	8966 (34,0%
Fair	17 (15,0%)	9409 (35,7%)
Poor	14 (12,4%)	3513 (13,3%)
Very poor	2 (1,8%)	824 (3,1%
Missing	22 (19,5%)	1774 (6,7%)
5. Over the last 12 months prior to your current hospita		-
how many times have you seen a doctor?	3 [2-7]	5 [2-10
how many times have you been admitted to the hospital (Emergency room, any ward)?	1 [1-3]	1 [0-2

	0[= 1]	
how many times have you been admitted to the hospital (Emergency room, any ward)?	1 [1-3]	



how many nights in total have you spent in hospital?	3 [1-10]	5 [0-15]
	L 1	

6. How many different medications do you take routinely ea	ch day (prior to hospit	alisation)?
1-2	20 (17,7%)	4936 (18,7%)
3-5	23 (20,4%)	6672 (25,3%)
More than 5	10 (8,8%)	7906 (30,0%)
None	30 (26,5%)	3926 (14,9%)
I do not know	7 (6,2%)	1077 (4,1%)
Missing	23 (20,4%)	1823 (6,9%)
7. Do you have health insurance?		
Yes, private insurance only	5 (4,4%)	4253 (16,1%)
Yes, public insurance only	3 (2,7%)	12350 (46,9%)
Yes, both	1 (0,88%)	3229 (12,3%)
None	74 (65,5%)	2938 (11,2%)
I prefer not to answer	4 (3,5%)	1314 (5,0%)
Missing	26 (23,0%)	2256 (8,6%)
8. What was your weight 5 years ago?	61 [55-75]	72 [60-85]
I do not know	40 (35,4%)	6267 (23,8%)
9a. Have you lost weight within the last 3 months? Yes, intentionally	11 (9,7%)	2468 (9,4%)
Yes, unintentionally	29 (25,7%)	10065 (38,2%)
No, my weight stayed the same	23 (20,4%)	7869 (29,9%)
No, I gained weight	33 (29,2%)	2857 (10,8%)
I do not know	12 (10,6%)	2512 (9,5%)
Missing	5 (4,4%)	569 (2,2%)
9b. If yes, how many kg did you lose?	8 [4-14]	6 [4-10]
I do not know	7 (17,5%)	1997 (15,9%)
10. Did you know about your hospitalisation two days before admission? (Yes)	30 (38,0%)	9602 (40,6%)
11. Please indicate if you		
were weighed at admission	49 (54,4%)	14112 (57,4%)
were informed about your nutrition status	33 (36,7%)	8364 (34,3%)
were informed about nutrition care options	19 (21,1%)	7769 (31,9%)
received special nutrition care	21 (23,6%)	7463 (30,7%)
12. How well have you eaten in the week before you were a	dmitted to the hospital	?

12. How were have you catch in the week before you were dumited to the hospital.		
More than normal	1 (0,89%)	1163 (4,4%)
Normal	51 (45,5%)	15832 (60,4%)

About 3/4 of normal	15 (13,4%)	2728 (10,4%)
About half of normal	31 (27,7%)	3287 (12,5%)
About a quarter to nearly nothing	12 (10,7%)	2496 (9,5%)
I do not know	1 (0,89%)	2490 (9,3%) 264 (1,0%)
	1 (0,89%)	443 (1,7%)
Missing	1 (0,09%)	443 (1,7%)
13. In general, how satisfied are you with the food at the ho	ospital?	
Very satisfied	21 (18,8%)	6066 (23,1%)
Somewhat satisfied	34 (30,4%)	8166 (31,2%)
Neutral	25 (22,3%)	4867 (18,6%)
Dissatisfied	5 (4,5%)	2103 (8,0%)
Very dissatisfied	2 (1,8%)	726 (2,8%)
I do not know	3 (2,7%)	2021 (7,7%)
Missing	22 (19,6%)	2264 (8,6%)
	(,	(0,0,0)
14. Did you get any help with eating TODAY?		
Yes, from family or friends	16 (14,3%)	2448 (9,3%)
Yes, from hospital staff	16 (14,3%)	1670 (6,4%)
No	55 (49,1%)	19193 (73,2%)
I do not know	-	302 (1,2%)
Missing	25 (22,3%)	2600 (9,9%)
15. Were you able to eat without interruption TODAY? (Yes)	68 (80,0%)	17231 (74,2%)
16a. Please indicate how much hospital food you ate for lu	nch or dinner TODAY	
About all	28 (25,0%)	12095 (46,1%)
1/2	47 (42,0%)	6486 (24,7%)
1/4	26 (23,2%)	3353 (12,8%)
Nothing	9 (8,0%)	3130 (11,9%)
Missing	2 (1,8%)	1149 (4,4%)
Nicol y	2 (1,070)	1110 (1,170)
16b. The portion size of the meal I ordered TODAY was		
Standard	32 (28,6%)	15979 (61,0%)
Smaller	7 (6,3%)	2367 (9,0%)
Larger	4 (3,6%)	1119 (4,3%)
I do not know	19 (17,0%)	2429 (9,3%)
Missing	50 (44,6%)	4319 (16,5%)
17. If you did not eat everything of your meal, please tell us	s why:	
I did not like the type of food offered	18 (22,0%)	2061 (15,9%)
I did not like the smell/taste of the food	6 (7,3%)	1569 (12,1%)
The food did not fit my cultural/religious preferences	-	143 (1,1%)
The food was too hot	-	59 (0,45%)



The food was too cold	4 (4,9%)	296 (2,3%)
Due to food allergy/intolerance	-	96 (0,74%)
I was not hungry at that time	12 (14,6%)	2205 (17,0%)
I do not have my usual appetite	33 (40,2%)	3830 (29,5%)
I have problems chewing/swallowing	4 (4,9%)	761 (5,9%)
I normally eat less than what was served	6 (7,3%)	1427 (11,0%)
I had nausea/vomiting	7 (8,5%)	1029 (7,9%)
I was too tired	21 (25,6%)	789 (6,1%)
I cannot eat without help	1 (1,2%)	208 (1,6%)
I was not allowed to eat	6 (7,3%)	1289 (9,9%)
I had an exam, surgery, or test and missed my meal	-	766 (5,9%)
I did not get requested food	-	139 (1,1%)
No answer given	3 (3,7%)	1829 (14,1%)

18. Enter the number of glasses/cups of the drinks you co	onsumed in the last 24 hou	rs
Water	4 [2-6]	3 [2-5]
Теа	1 [1-2]	1 [0-2]
Coffee	1 [0-1]	1 [1-2]
Milk	1 [1-2]	1 [0-1]
Fruit juice	1 [1-2]	1 [0-2]
Soft drinks	0 [0-0]	0 [0-1]
Nutrition drink	1 [0-1]	0 [0-1]
Other	1 [0-1]	0 [0-1]
19a. Did you eat any food apart from hospital food TODAY?	26 (35,6%)	6580 (28,9%)
19b. If yes, what did you eat?		
Sweet snacks	2 (7,7%)	1928 (29,3%)
Salty snacks	3 (11,5%)	749 (11,4%)
Homemade food	6 (23,1%)	963 (14,6%)
Fruits	5 (19,2%)	2551 (38,8%)
Dairy products	4 (15,4%)	724 (11,0%)
Food delivered/restaurant	3 (11,5%)	293 (4,5%)
Sandwich	9 (34,6%)	421 (6,4%)
Other	1 (3,8%)	1042 (15,8%)
20. How has your food intake changed since your hospita	l admission?	

Increased	4 (3,6%)	3706 (14,1%)
Decreased	43 (38,4%)	7774 (29,7%)
Stayed the same	26 (23,2%)	10331 (39,4%)
I do not know	6 (5,4%)	1690 (6,4%)
Missing	33 (29,5%)	2712 (10,3%)



21. TODAY I feel		
Stronger than at admission	42 (37,5%)	11240 (42,9%)
Weaker than at admission	23 (20,5%)	4893 (18,7%)
Same as at admission	20 (17,9%)	7428 (28,3%)
I was admitted today	4 (3,6%)	662 (2,5%)
I do not know	18 (16,1%)	1384 (5,3%)
Missing	5 (4,5%)	606 (2,3%)
22. Can you walk without assistance TODAY?		
Yes	82 (73,2%)	15438 (58,9%)
No, only with assistance	15 (13,4%)	6458 (24,6%)
No, I stay in bed	14 (12,5%)	3018 (11,5%)
Missing	1 (0,89%)	1299 (5,0%)
23. Did anyone help you complete this questionnaire?	39 (44,3%)	15731 (65,1%)

